

Algebra I

LSSM – Algebra I		Explicit Component(s) of Rigor		
Code	Standard	Conceptual Understanding	Procedural Skill and Fluency	Application
A1: N-RN.B.3	Explain why the sum or product of two rational numbers is rational; that the sum of a rational number and an irrational number is irrational; and that the product of a nonzero rational number and an irrational number is irrational.	✓		
A1: N-Q.A.1	Use <u>units</u> as a way to <u>understand</u> problems and to guide the solution of multi-step problems; <u>choose and interpret</u> units consistently in formulas; <u>choose and interpret</u> the scale and the origin in graphs and data displays. *	✓		
A1: N-Q.A.2	<u>Define</u> appropriate quantities for the purpose of descriptive modeling. *	✓		
A1: N-Q.A.3	<u>Choose</u> a level of accuracy appropriate to limitations on measurement when reporting quantities. *	✓		
A1: A-SSE.A.1	<u>Interpret</u> expressions that represent a quantity in terms of its context. *	✓		
A1: A-SSE.A.1a	<u>Interpret</u> parts of an expression, such as terms, factors, and coefficients. *	✓		
A1: A-SSE.A.1b	<u>Interpret</u> complicated expressions by viewing one or more of their parts as a single entity. <i>For example, interpret $P(1+r)^n$ as the product of P and a factor not depending on P.</i> *	✓		
A1: A-SSE.A.2	Use the structure of an expression to <u>identify</u> ways to rewrite it. <i>For example, see $x^4 - y^4$ as $(x^2)^2 - (y^2)^2$, or see $2x^2 + 8x$ as $(2x)(x) + 2x(4)$, thus recognizing it as a polynomial whose terms are products of monomials and the polynomial can be factored as $2x(x+4)$.</i>	✓	✓	
A1: A-SSE.B.3	<u>Choose and produce</u> an equivalent form of an expression to <u>reveal and explain</u> properties of the quantity represented by the expression. *	✓	✓	
A1: A-SSE.B.3a	<u>Factor</u> a quadratic expression to <u>reveal</u> the zeros of the function it defines. *	✓	✓	
A1: A-SSE.B.3b	<u>Complete the square</u> in a quadratic expression to <u>reveal</u> the maximum or minimum value of the function it defines. *	✓	✓	
A1: A-SSE.B.3c	Use the properties of exponents to <u>transform</u> expressions for exponential functions emphasizing integer exponents. <i>For example, the growth of bacteria can be modeled by either $f(t) = 3^{(t+2)}$ or $g(t) = 9(3^t)$ because the expression $3^{(t+2)}$ can be rewritten as $(3^t)(3^2) = 9(3^t)$.</i> *		✓	
A1: A-APR.A.1	<u>Understand</u> that polynomials form a system analogous to the integers, namely, they are closed under the operations of addition, subtraction, and multiplication; <u>add, subtract, and multiply</u> polynomials.	✓	✓	
A1: A-APR.B.3	<u>Identify</u> zeros of quadratic functions, and <u>use the zeros</u> to <u>sketch</u> a graph of the function defined by the polynomial.	✓	✓	

A1: A-CED.A.1	Create equations and inequalities in one variable and use them to solve problems. Include equations arising from linear, quadratic, and exponential functions.*	✓	✓	✓
A1: A-CED.A.2	Create equations in two or more variables to represent relationships between quantities; graph equations on coordinate axes with labels and scales.*	✓	✓	
A1: A-CED.A.3	Represent constraints by equations or inequalities, and by systems of equations and/or inequalities, and interpret solutions as viable or nonviable options in a modeling context. For example, represent inequalities describing nutritional and cost constraints on combinations of different foods.*	✓		✓
A1: A-CED.A.4	Rearrange formulas to highlight a quantity of interest, using the same reasoning as in solving equations. For example, rearrange Ohm's law $V = IR$ to highlight resistance R .*		✓	
A1: A-REI.A.1	Explain each step in solving a simple equation as following from the equality of numbers asserted at the previous step, starting from the assumption that the original equation has a solution. Construct a viable argument to justify a solution method.	✓		
A1: A-REI.B.3	Solve linear equations and inequalities in one variable, including equations with coefficients represented by letters.		✓	
A1: A-REI.B.4	Solve quadratic equations in one variable.		✓	
A1: A-REI.B.4a	Use the method of completing the square to transform any quadratic equation in x into an equation of the form $(x - p)^2 = q$ that has the same solutions. Derive the quadratic formula from this form.		✓	
A1: A-REI.B.4b	Solve quadratic equations by inspection (e.g., for $x^2 = 49$), taking square roots, completing the square, the quadratic formula and factoring, as appropriate to the initial form of the equation. Recognize when the quadratic formula gives complex solutions and write them as "no real solution".*	✓	✓	
A1: A-REI.C.5	Prove that, given a system of two equations in two variables, replacing one equation by the sum of that equation and a multiple of the other produces a system with the same solutions.	✓		
A1: A-REI.C.6	Solve systems of linear equations exactly and approximately (e.g., with graphs), focusing on pairs of linear equations in two variables.		✓	
A1: A-REI.D.10	Understand that the graph of an equation in two variables is the set of all its solutions plotted in the coordinate plane, often forming a curve (which could be a line).	✓		
A1: A-REI.D.11	Explain why the x -coordinates of the points where the graphs of the equations $y = f(x)$ and $y = g(x)$ intersect are the solutions of the equation $f(x) = g(x)$; find the solutions approximately, e.g., using technology to graph the functions, make tables of values, or find successive approximations. Include cases where $f(x)$ and/or $g(x)$ are linear, polynomial, rational, piecewise linear (to include absolute value), and exponential functions.*	✓	✓	
A1: A-REI.D.12	Graph the solutions to a linear inequality in two variables as a half-plane (excluding the boundary in the case of a strict inequality), and graph the solution set to a system of linear inequalities in two variables as the intersection of the corresponding half-planes.		✓	

A1: F-IF.A.1	<u>Understand</u> that a function from one set (called the domain) to another set (called the range) assigns to each element of the domain exactly one element of the range. If f is a function and x is an element of its domain, then $f(x)$ denotes the output of f corresponding to the input x . The graph of f is the graph of the equation $y = f(x)$.	✓		
A1: F-IF.A.2	<u>Use</u> function notation, <u>evaluate</u> functions for inputs in their domains, and <u>interpret</u> statements that use function notation in terms of a context.	✓	✓	
A1: F-IF.A.3	<u>Recognize</u> that sequences are functions whose domain is a subset of the integers. <u>Relate</u> arithmetic sequences to linear functions and geometric sequences to exponential functions.	✓		
A1: F-IF.B.4	For linear, piecewise linear (to include absolute value), quadratic, and exponential functions that model a relationship between two quantities, <u>interpret</u> key features of graphs and tables in terms of the quantities, and <u>sketch</u> graphs showing key features given a verbal description of the relationship. <i>Key features include: intercepts; intervals where the function is increasing, decreasing, positive, or negative; relative maximums and minimums; symmetries; and end behavior.</i> *	✓		
A1: F-IF.B.5	<u>Relate</u> the domain of a function to its graph and, where applicable, to the quantitative relationship it describes. <i>For example, if the function $h(n)$ gives the number of person-hours it takes to assemble n engines in a factory, then the positive integers would be an appropriate domain for the function.</i> *	✓		
A1: F-IF.B.6	<u>Calculate and interpret</u> the average rate of change of a linear, quadratic, piecewise linear (to include absolute value), and exponential function (presented symbolically or as a table) over a specified interval. <u>Estimate</u> the rate of change from a graph.*	✓	✓	
A1: F-IF.C.7	<u>Graph</u> functions expressed symbolically and <u>show</u> key features of the graph, by hand in simple cases and using technology for more complicated cases.*	✓	✓	
A1: F-IF.C.7a	<u>Graph</u> linear and quadratic functions and <u>show</u> intercepts, maxima, and minima.*	✓	✓	
A1: F-IF.C.7b	<u>Graph</u> piecewise linear (to include absolute value) and exponential functions.*		✓	
A1: F-IF.C.8	<u>Write</u> a function defined by an expression in different but equivalent forms to <u>reveal and explain</u> different properties of the function.	✓	✓	
A1: F-IF.C.8a	<u>Use the process of factoring and completing the square</u> in a quadratic function to <u>show</u> zeros, extreme values, and symmetry of the graph, and <u>interpret</u> these in terms of a context.	✓	✓	
A1: F-IF.C.9	<u>Compare</u> properties of two functions (linear, quadratic, piecewise linear [to include absolute value] or exponential) each represented in a different way (algebraically, graphically, numerically in tables, or by verbal descriptions). <i>For example, given a graph of one quadratic function and an algebraic expression for another, determine which has the larger maximum.</i>	✓	✓	
A1: F-BF.A.1	<u>Write</u> a linear, quadratic, or exponential function that describes a relationship between two quantities.*	✓	✓	
A1: F-BF.A.1a	<u>Determine</u> an explicit expression, a recursive process, or steps for calculation from a context.*	✓	✓	

A1: F-BF.B.3	Identify the effect on the graph of replacing $f(x)$ by $f(x) + k$, $k f(x)$, $f(kx)$, and $f(x + k)$ for specific values of k (both positive and negative). Without technology, <u>find</u> the value of k given the graphs of linear and quadratic functions. With technology, <u>experiment</u> with cases and <u>illustrate an explanation</u> of the effects on the graph that include cases where $f(x)$ is a linear, quadratic, piecewise linear (to include absolute value) or exponential function.	✓	✓	
A1: F-LE.A.1	<u>Distinguish</u> between situations that can be modeled with linear functions and with exponential functions.*	✓		
A1: F-LE.A.1a	<u>Prove</u> that linear functions grow by equal differences over equal intervals, and that exponential functions grow by equal factors over equal intervals.*	✓		
A1: F-LE.A.1b	<u>Recognize</u> situations in which one quantity changes at a constant rate per unit interval relative to another.*	✓		
A1: F-LE.A.1c	<u>Recognize</u> situations in which a quantity grows or decays by a constant percent rate per unit interval relative to another.*	✓		
A1: F-LE.A.2	<u>Construct</u> linear and exponential functions, including arithmetic and geometric sequences, given a graph, a description of a relationship, or two input-output pairs (include reading these from a table).*	✓	✓	
A1: F-LE.A.3	<u>Observe</u> using graphs and tables that a quantity increasing exponentially eventually exceeds a quantity increasing linearly, quadratically, or (more generally) as a polynomial function.*	✓		
A1: F-LE.B.5	<u>Interpret</u> the parameters in a linear or exponential function in terms of a context.*	✓		
A1: S-ID.A.2	<u>Use statistics</u> appropriate to the shape of the data distribution to <u>compare</u> center (median, mean) and spread (interquartile range, standard deviation) of two or more different data sets.*	✓	✓	
A1: S-ID.A.3	<u>Interpret</u> differences in shape, center, and spread in the context of the data sets, accounting for possible effects of extreme data points (outliers).*	✓		
A1: S-ID.B.5	<u>Summarize</u> categorical data for two categories in two-way frequency tables. <u>Interpret</u> relative frequencies in the context of the data (including joint, marginal, and conditional relative frequencies). <u>Recognize</u> possible associations and trends in the data.*	✓	✓	
A1: S-ID.B.6	<u>Represent</u> data on two quantitative variables on a scatter plot, and <u>describe</u> how the variables are related.*	✓	✓	
A1: S-ID.B.6a	<u>Fit</u> a function to the data; <u>use functions</u> fitted to data to <u>solve problems</u> in the context of the data. <u>Use given functions or choose a function</u> suggested by the context. Emphasize linear and quadratic models.*	✓	✓	✓
A1: S-ID.B.6b	<u>Informally assess</u> the fit of a function <u>by plotting and analyzing</u> residuals.*	✓	✓	
A1: S-ID.B.6c	<u>Fit</u> a linear function for a scatter plot that suggests a linear association.*		✓	
A1: S-ID.C.7	<u>Interpret</u> the slope (rate of change) and the intercept (constant term) of a linear model in the context of the data.*	✓		

A1: S-ID.C.8	<u>Compute</u> (using technology) and <u>interpret</u> the correlation coefficient of a linear fit. *	✓	✓	
A1: S-ID.C.9	<u>Distinguish</u> between correlation and causation. *	✓		

*Modeling standard